

Lipid profile reference intervals in individuals from Maracaibo, Venezuela: an insight from the Maracaibo City Metabolic Syndrome prevalence study

Valmore J. Bermúdez, MD, MPH, PhD*¹, Luis Miguel Bello, BSc¹, Ariana Naguib, BSc¹, Roberto J. Añez, BSc¹, Yanalith Fortul, BSc¹, Alexandra Toledo, BSc¹, Juan J. Salazar, BSc¹, Yaquelin Torres, BSc¹, Verónica Angulo, MD¹, Carlos Silva Paredes, MD¹, Sergia Linares, MgSc¹, Nailet Arraiz, PhD¹, Carem Prieto, MgSc¹, Enrique Pacheco, MD¹, Maricarmen Chacín, MD¹, Joselyn Rojas, MD, MSc^{1,2}, Miguel Aguirre, MD, MSc^{1,3}, Marjorie Villalobos, MD¹, Climaco Cano Ponce, PharmD¹, Rafael París Marciano, MD, PhD

¹Endocrine and Metabolic Diseases Research Center, The University of Zulia, Maracaibo –Venezuela.

²Institute of Clinical Immunology. University of Los Andes. Mérida –Venezuela.

³Endocrinology Unit, I.A.H.U.L.A, Mérida –Venezuela.

*Correspondence Author: Valmore J. Bermúdez, MD, MPH, PhD. Endocrine and Metabolic Diseases Research Center, The University of Zulia, Maracaibo – Venezuela.

E-mail: valmore@gmail.com

This paper is part of the academic duties of Dr. Valmore Bermúdez within the Master in Advanced Endocrinology at the University of Alcalá de Henares.

Director Dr. Melchor Álvarez de Mon Soto

Recibido: 16/04/2012 Aceptado: 20/06/2012

24

Abstract

Objectives: Cardiovascular diseases are the main cause of death in adults worldwide. Dyslipidemias are an important risk factor, which is why committees like the ATP III have established cut-off points for lipid profiles dichotomizing the diagnosis instead of setting normal intervals. Currently, there is no general consensus about the reference intervals in our population, so the purpose of this paper is to establish lipid profile reference ranges in Maracaibo City, Venezuela.

Materials and Methods: A cross-sectional study was made, enrolling 2,043 randomly selected individuals from both genders over 18 years old, for the Maracaibo City Metabolic Syndrome Prevalence Study (MMSPS). To determine the reference population, the patients with pathologic history and medication intake that could modify the lipid profile were taken out. Extreme values were calculated mathematically for each lipid variable and then excluded. The results were expressed in absolute and relative frequencies, medians and 25th-75th percentiles for the general and gender classified population.

Results: The medians and percentiles for the reference population (n=434) were indicated respectively for women (n=221; 50.9%) and men (n=213; 49.1%). For HDL-C (mg/dL): 48.00 (42.00-56.00) and 43.00 (37.00-50.00); Triglycerides (mg/dL): 70.98 (50.35-102.53) and 78.50 (57.00-126.63); VLDL-C (mg/dL): 14.20 (10.07-20.48) and 15.70 (11.40-25.33). Were raised general reference values for Total Cholesterol (mg/dL): 176.00 (149.50-201.50); LDL-C (mg/dL): 110.23 (86.25-132.80) and Lp(a) (mg/dL): 25.20 (18.40-33.85).

Key Words: Reference Intervals, Lipid Profile, Cardiovascular Disease, Dyslipidemia.

Introducción

Cardiovascular diseases are the primary cause of death worldwide¹, and Venezuela is no exception to this reality as this pathology stands as the first of the 25 leading causes of death in both men and woman and among individuals 65 -74 years in the country². However, the fact that a high proportion of vulnerable individuals are young adults helps to understand the importance of defining the population at risk of suffering from cardiovascular disease in a timely manner³.

Dyslipidemia constitutes the fundamental factor for atherogenesis and is considered one of the most important cardiovascular risk factors⁴⁻⁶, being directly related to modifiable factors such as diet or lifestyle habits, as well as non-modifiable factors from each individual like genetic predisposition⁷⁻⁹. For this reason, various committees including the Adult Treatment Panel III (ATP III) have defined and established reference values for plasma lipids⁸ in order to reduce the risk of developing cardiovascular disease¹⁰. Taking into account that lifestyle, genetics, and environmental conditions vary in each population, countries such as China, Spain and Turkey have conducted studies to determine their own cut-off values¹¹⁻¹³ for objectively establishing abnormal lipid profiles that could be considered a cardiovascular risk for their population. Given these variations in lipid determination and the diversity of factors that influence the behavior of dyslipidemia in each population, it is necessary to determine lipid profile reference intervals in order to amplify the spectrum of evaluation that provides the dichotomization of a specific diag-

nosis. Proposals such as this have also been designed in countries like India, Mexico and Colombia¹⁴⁻¹⁹.

Therefore, taking into consideration that mortality rate from cardiovascular events in the Zulia population is among the highest in Venezuela, the lack of epidemiological analyzing resident lipid behavior, and whether or not the values assigned by the ATP III committee are useful to assess the risk in the local population, it's of extreme importance to evaluate the presence of dyslipidemia in the Maracaibo Municipality, Zulia State, and determine new and appropriate local reference intervals for plasma lipids.

Ethical considerations

Individuals were included in the study after signing a written consent form prior to obtaining a history and physical examination. The study design was approved by the Ethics Committee of The Metabolic and Endocrine Diseases Research Center "Dr. Félix Gómez".

Selection of Individuals

The sample method was already published in the MMSPS cross-sectional proposal (20), yet the main features will be mentioned. Using population estimations for the population of Maracaibo (1,428,043 for 2007 according to the National Institute of Statistics), the sample size estimate was calculated to be 1,986 individuals with or above 18 years of age. Moreover, taking into account that in a previous pilot study approximately 10% of the subjects rejected being part of the study (unpublished data), an oversampling number was calculated (200 individuals); the overall number of individuals was 2,230 (with 244 subjects - 12.0% - added because of the oversampling method) were randomly selected between July 2008 and July 2011²⁰. The only inclusion criterion was to have ≥ 18 years of age. The city of Maracaibo is divided into parishes and each of these was proportionally sampled: Antonio Borjas Romero, Bolívar, Cacique Mara, Caracciolo Parra Pérez, Cecilio Acosta, Cristo de Aranza, Coquivacoa, Chiquinquirá, Francisco Eugenio Bustamante, Idelfonso Vásquez, Juana de Ávila, Luis Hurtado Higuera, Manuel Dagnino, Olegario Villalobos, Raúl Leoni, Santa Lucía, San Isidro, and Venancio Pulgar. The sampling process was undertaken using a 2-phase method: During the first phase, the sorting was random and stratified—where each stratum was represented by sectors from each of the 18 parishes—choosing 4 from each parish. The second sampling was stratified to represent a city block, in which they were selected using a random number generation tool. Once the houses were selected, every adult in the family unit from the selected city blocks was invited to participate in the study and were interviewed on prior written consent, and subjected to a routine medical examination using the clinical chart provided by the Health and Social Development Ministry of Venezuela as data collecting tool. The sample used in this

research was of 2,043 individuals that gathered all the required biochemical parameters for the study.

Anthropometric assessment

Body Mass Index (BMI): Weight was determined using a digital scale (Tanita, TBF-310 Body Composition Analyzer GS, Tokyo - Japan), while height was obtained using a vertical strip calibrated in centimeters and millimeters. The patients were barefooted and in light clothing at the time of the measurements. BMI was calculated by the mathematical formula: $\text{Weight (kg)} / \text{Height}^2 (\text{m}^2)$.

Laboratory tests

Prior 8 to 12 hours of fasting, 5 cm³ of blood was extracted from each individual by venipuncture of the antecubital vein, being placed afterwards in test tubes and centrifuged at 4000 rpm for 10 minutes; serum was extracted and placed in polypropylene test tubes for subsequent freezing at -70 °C. The time between sample collection and processing did not exceed three months. Total Cholesterol was estimated by enzymatic colorimetric methods (Wiener Lab SAIC). Triglycerides (TAG) and High Density Lipoprotein (HDL-C) were evaluated using an enzyme-colorimetric commercial kit (Human Gesellschaft Biochemica and Diagnostica MBH) following the manufacturer instruction. Low Density Lipoprotein (LDL-C) was estimated using the Friedewald formula²¹ in subjects with triglycerides below 400 mg/dl¹⁵. Those who obtained TAG > 400 mg/dl, underwent measurement of their concentration by electrophoresis techniques. Very Low Density Lipoprotein was calculated indirectly from the mathematical expression $[\text{TAG}/5]$. Lipoprotein (a) [Lp (a)] was estimated through the latex turbidimetric method, Human Gesellschaft für Biochemica und Diagnostica MBH, Germany. In this method, the presence of Lp(a) in the sample causes agglutination of latex particles coated with antibodies against Lp(a). The agglutination is proportional to the Lp(a) concentration in the sample and can be measured by turbidimetry (22). For glucose determination a colorimetric enzymatic glucose oxidase kit (Sigma, USA) was used, using both standard and 0.02 ml of sample, then adding 2 ml of the enzyme reagent. Incubated for 10 minutes at 20-25 °C or 5 minutes at 37 °C, then absorption was measured at 500nm against a target reagent. Basal Insulin plasma levels were quantified through the international DRG insulin kit. Inc. USA. New Jersey. A solid phase immunoassay of two sites that uses two monoclonal antibodies directed to two antigenic determinants on different sites in the insulin molecule which are detected by reaction with 3,3', 5,5'-tetramethylbenzidine and the addition of an acid to stop the reaction and provide a colorimetric endpoint read by spectrophotometer. Detection limit <1 mU/l. Calculation of insulin resistance was conducted applying the HOMA-Calculator, Homa-IR2, which is an update and adaptation by Jonathan Levy et al.²³.

Exclusion Criteria

For the determination of reference intervals in biological variables such as lipid profile is necessary to establish a reference population²⁵⁻²⁸, for which exclusion criteria listed in Table 1 was used.

Table 1: Exclusion criteria based on clinical history and laboratory tests to define the reference individuals.

Exclusion Criteria	
Obesity as BMI (≥ 30 kg/m ²)	Personal history of Hypertension
Glycemia ≥ 126 mg/dL	Personal history of Diabetes Mellitus
HOMA-IR ≥ 2	Personal history of Thyroid and/or Hepatic disease
Acanthosis Nigricans	Personal history of Polycystic Ovary Syndrome
Alcohol Consumption	Personal history of Acute Myocardial Infarction
Consumption of medications that modify lipid profile and/or Alcohol	Personal history of Angina. Arrhythmia and/or Cerebrovascular disease

Calculation of Extreme Values

Once the reference population was established, we proceeded to calculate the extreme values for each lipid variable to avoid any bias that might have an influence on the reference intervals distribution^{24,29,30}. This was performed using the mathematical expression:

$$[\text{Upper limit: } Q_3 + 1,5 (Q_3 - Q_1)]$$

$$[\text{Lower limit: } Q_1 - 1,5 (Q_3 - Q_1)]$$

Where:

Q_1 = firstquartile (25th percentile)

Q_3 = thirdquartile (75th percentile)

Therefore any result above the upper limit or below the lower limit, is an extreme value³⁰ and was removed from the study^{24,29}.

Statistical Analysis

Statistical analysis was performed using the Statistical Package for Social Sciences for Windows, version 19.0 (SPSS Inc.) and the R-project program, version 2.11.0. The qualitative variables (Gender and Age group) were expressed as absolute and relative frequencies. The quantitative variables: Total Cholesterol, Triglycerides, VLDL-C, LDL-C, HDL-C and Lp(a), were tested by goodness of fit (normality and homoscedasticity). To check the normality of the variables the Geary test was used. We applied the Levene test to evaluate the homoscedasticity. For those who showed a normal distribution results were expressed as mean \pm SD (standard deviation) with their respective confidence intervals at 95% (CI 95%). The differences between these were established using the Student's Ttest or one-way ANOVA with the Tukey post-hoc test depending on the case, considering $p < 0.05$ as statistically significant. The results of abnormally distributed

variables were expressed as medians and percentiles p2, 5, p5, p10, p25, p50, p75, p90, p95, p97, 5. After statistical processing of the variables, the reference values for the lipid profile were determined using p25 as lower value and p75 as upper value. The differences between these were calculated using the Mann-Whitney U test or ANOVA with Bonferroni correction when indicated.

Results

General Characteristics of the Population

The Maracaibo City Metabolic Syndrome Prevalence Study (MMSPS) has an overall sample of 2,043 individuals of both genders, male: $n=952$ (46.6%), female: $n=1,091$ (53.4%). The mean population age was 39.78 ± 15.45 (CI 95%: 39.78 to 40.45) years. By stratifying the age variable there was a predominance of the age group 20-29 years with 24.6% (Table 2), the group of 40 to 49, 21.1%; the group of 30 to 39 years with 17.7%, so it was observed that about 63.4% of the population was aged between 20 and 49 years. The population behavior for each of the lipid profile variables by gender are shown in Table 3.

Table 2: Distribution of general population by age group and gender. Maracaibo Municipality, Zulia State, 2011

Age Group (Years)	Female		Male		Total	
	n	(%)	n	(%)	n	(%)
18 – 19	96	(8.8%)	71	(7.5%)	167	(8.2%)
20 – 29	233	(21.4%)	270	(28.4%)	503	(24.6%)
30 – 39	183	(16.8%)	178	(18.7%)	361	(17.7%)
40 – 49	247	(22.6%)	184	(19.3%)	431	(21.1%)
50 – 59	189	(17.3%)	160	(16.8%)	349	(17.1%)
≥ 60	143	(13.1%)	89	(9.3%)	232	(11.4%)
Total	1091	(100%)	952	(100%)	2043	(100%)

Table 3: Percentiles and measures of central tendency and dispersion of the lipid profile in the general population of the Maracaibo Municipality, Zulia State, 2011.

Lipid Profile (mg/dL)	Gender	Percentiles				
		Mean	SD	P25	P50	P75
HDL-C	Female	47.07	11.89	39.00	46.00	54.00
	Male	41.07	11.58	34.00	39.00	46.00
	Total	44.27	12.12	36.00	43.00	51.00
TAG	Female	116.90	85.30	64.00	96.00	143.00
	Male	146.08	117.37	76.00	117.00	174.00
	Total	130.50	102.53	68.82	105.00	159.40
VLDL	Female	23.38	17.06	12.80	19.20	28.60
	Male	29.22	23.47	15.20	23.40	34.80
	Total	26.10	20.50	13.76	21.00	31.88
Total Cholesterol	Female	194.25	44.67	162.00	190.00	220.00
	Male	189.33	47.36	159.00	184.00	214.00
	Total	191.96	46.00	161.00	187.00	217.00
LDL	Female	123.85	38.11	96.58	120.94	146.15
	Male	120.03	38.65	93.40	116.80	144.00
	Total	122.10	38.40	95.45	119.27	145.66
Lp(a)	Female	28.54	14.07	19.60	26.60	35.70
	Male	27.33	14.11	18.20	25.25	35.40
	Total	28.01	14.10	19.05	26.10	35.60

SD: Standard deviation

After applying the exclusion criteria, a population consisting of 434 healthy individuals, of whom 50.9% (n=221) were female and 49.1% (n= 213) were male was obtained. The meanaverage age of this group was 33.26 ± 14.10 years (CI 95% 31.93 to 34.59 years).In order to evaluate the lipid profile in the reference population, ex-

treme values were calculated to exclude them from the analysis, obtaining reference samples for each lipid variable (Table 4). Based on these reference samples, results obtained from analysis of each lipid profile variable are expressed in Table 5.

Table 4: Reference samples for lipid profile study grouped by gender and age group. Maracaibo Municipality, Zulia State, 2011.

Lipid Profile (mg/dL)	Gender	Age group (Years)						Total
		18-19	20-29	30-39	40-49	50-59	≥ 60	
		n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	
HDL-C	Female	35 (16.3%)	66 (30.7%)	45 (20.9%)	43 (20.0%)	16 (7.4%)	10 (4.7%)	215 (100%)
	Male	33 (15.9%)	82 (39.6%)	38 (18.4%)	15 (7.2%)	23 (11.1%)	16 (7.7%)	207 (100%)
	Total	68 (16.1%)	148 (35.1%)	83 (19.7%)	58 (13.7%)	39 (9.2%)	26 (6.2%)	422 (100%)
Triglycerides VLDL-C	Female	34 (16.0%)	63 (29.7%)	46 (21.7%)	43 (20.3%)	17 (8.0%)	9 (4.2%)	212 (100%)
	Male	32 (15.8%)	84 (41.6%)	38 (18.8%)	15 (7.4%)	18 (8.9%)	15 (7.4%)	202 (100%)
	Total	68 (15.9%)	147 (35.5%)	84 (20.3%)	58 (14.0%)	35 (8.5%)	24 (5.8%)	414 (100%)
LDL-C	Female	36 (17.1%)	63 (29.9%)	46 (21.8%)	43 (20.4%)	14 (6.6%)	9 (4.3%)	211 (100%)
	Male	33 (15.6%)	85 (40.3%)	37 (17.9%)	17 (8.1%)	23 (10.9%)	16 (7.6%)	211 (100%)
	Total	69 (16.4%)	148 (36.1%)	83 (19.7%)	60 (14.2%)	37 (8.8%)	25 (5.9%)	422 (100%)
Total Cholesterol	Female	36 (16.7%)	65 (30.2%)	46 (21.4%)	43 (20.0%)	15 (7.0%)	10 (4.7%)	215 (100%)
	Male	33 (15.7%)	85 (40.5%)	37 (17.6%)	17 (8.1%)	22 (10.5%)	16 (7.6%)	210 (100%)
	Total	69 (16.2%)	150 (35.3%)	83 (19.5%)	60 (14.1%)	37 (8.7%)	26 (6.1%)	425 (100%)
Lp (a)	Female	28 (15.6%)	53 (29.4%)	40 (22.2%)	38 (21.1%)	13 (7.2%)	8 (4.4%)	180 (100%)
	Male	29 (19.5%)	52 (34.9%)	23 (15.4%)	13 (8.7%)	16 (10.7%)	16 (10.7%)	149 (100%)
	Total	57 (17.3%)	105 (31.9%)	63 (19.1%)	51 (15.5%)	29 (8.8%)	24 (7.3%)	329 (100%)

Table 5: Lipid profile percentiles in the reference sample distributed by gender. Maracaibo Municipality, Zulia State, 2011.

Percentiles	Female						Male						Total					
	Total Cholesterol (mg/dL)	TAG (mg/dL)	VLDL-C (mg/dL)	LDL-C (mg/dL)	HDL-C (mg/dL)	Lp(a) (mg/dL)	Total Cholesterol (mg/dL)	TAG (mg/dL)	VLDL-C (mg/dL)	LDL-C (mg/dL)	HDL-C (mg/dL)	Lp(a) (mg/dL)	Total Cholesterol (mg/dL)	TAG (mg/dL)	VLDL-C (mg/dL)	LDL-C (mg/dL)	HDL-C (mg/dL)	Lp(a) (mg/dL)
p2.5	121.40	31.00	6.20	56.6	30.40	2.70	107.55	33.00	6.6	45.16	30.00	1.66	114.95	31.38	6.28	53.64	30.00	2.06
p5	126.80	34.47	6.89	62.72	33.80	8.80	117.55	38.15	7.63	57.44	31.00	3.39	122.30	36.00	7.20	61.66	32.00	6.18
p10	133.60	40.30	6.06	71.64	36.00	12.71	127.10	45.16	9.03	68.68	33.00	9.80	130.60	42.00	8.40	70.40	35.00	12.60
p25	151.00	50.35	10.07	87.20	42.00	18.63	147.00	57.00	11.40	85.60	37.00	18.25	149.50	53.75	10.75	86.25	39.00	18.40
p50	176.00	70.98	14.20	109.60	48.00	25.35	175.00	78.50	15.70	110.80	43.00	24.90	176.00	74.69	14.94	110.23	45.00	25.20
p75	202.00	102.53	20.48	130.20	56.00	34.30	200.25	126.63	25.33	134.60	50.00	33.65	201.50	109.93	21.99	132.80	53.00	33.85
p90	231.40	133.16	26.63	150.85	63.40	40.67	234.60	173.70	34.74	159.52	61.00	41.80	231.40	154.90	30.98	157.54	62.00	41.10
p95	248.40	154.37	30.87	162.20	70.20	46.56	251.90	199.00	39.80	178.72	67.20	48.65	249.70	181.00	36.20	173.01	68.00	46.85
p97.5	271.00	170.75	34.15	176.14	75.60	50.84	269.73	207.93	41.59	195.82	76.40	51.93	270.00	199.00	39.80	185.41	75.43	50.98


HDL-C

The reference sample for the calculation of HDL-C was 422 individuals (Table 4) with anaverage age of 33.18 ± 14.15

years (CI 95% 31.83 to 34.53 years).In the overall sample a median of 45.00 (39.00 to 53.00) mg/dL was obtained. When divided by gender it showed that female individuals

had a median of 48.00 (42.00 to 56.00) mg/dL, while men obtained a value of 42.00 (37.00 to 50.00) mg/dL. In the analysis of both groups statistically significant differences were appreciated ($p=1.437 \times 10^{-6}$); (Figure 1, panel A). The median concentrations of HDL-C obtained by gathering age group by gender are given in Table 6 where statistically

Figure 1: Comparison of median values of HDL-C, VLDL-C and Lp(a) in a reference sample by gender. Maracaibo Municipality, Zulia State, 2011.



significant differences between men and women in age groups between 18-19; 30-39; 40-49 and in the 60 and over group are shown (Figure 3). No statistically significant differences were observed in different age groups when studying HDL-C concentrations by gender while gathering them by age group.

Figure 3: Comparison of medians of concentration of HDL-C in a reference sample by age group. Maracaibo Municipality, Zulia State, 2011.




Table 6: Comparison of reference intervals of Lipid profile distributed by gender and age group. Maracaibo Municipality, Zulia State, 2011.

Age Group (Years)		HDL-C (mg/dL)			TAG (mg/dL)			VLDL-C (mg/dL)			Total Cholesterol (mg/dL)			LDL-C (mg/dL)			Lp(a) (mg/dL)		
		P25	P50	P75	P25	P50	P75	P25	P50	P75	P25	P50	P75	P25	P50	P75	P25	P50	P75
18 – 19	Female	41.00	50.00	56.00	44.57	58.50	93.00	8.91	11.70	18.60	143.00	156.50	176.00	79.35	92.18	108.57	13.07	21.87	34.62
	Male	38.00	42.00	48.00	44.25	68.00	100.33	8.85	13.60	20.06	124.00	146.00	166.00	70.90	85.00	106.82	11.70	18.30	26.15
	<i>P^a</i>	0.033*			0.534			0.534			0.036*			0.175			0.196		
20 – 29	Female	41.75	49.00	58.25	49.00	65.00	82.00	9.80	13.00	16.40	136.00	167.00	189.00	77.40	100.50	120.40	18.10	22.80	31.90
	Male	40.00	45.00	52.00	51.00	67.01	106.75	10.20	13.40	21.35	144.50	172.00	193.00	79.70	107.60	126.50	18.40	23.95	32.72
	<i>P^a</i>	0.103			0.142			0.142			0.420			0.170			0.980		
30 – 39	Female	43.00	49.00	54.00	47.00	64.00	102.17	9.40	12.80	20.42	149.75	171.50	206.25	91.82	111.00	134.99	20.52	26.10	30.92
	Male	35.00	39.50	51.00	75.25	114.50	168.50	15.05	22.90	33.70	162.50	187.00	210.50	98.00	120.20	147.30	19.70	24.80	31.70
	<i>P^a</i>	0.006*			6.59x10 ⁻⁵ *			6.59x10 ⁻⁵ *			0.146			0.223			0.836		
40 – 49	Female	39.00	46.00	54.00	55.00	82.09	115.00	11.00	16.41	23.00	167.00	189.00	207.00	105.00	124.80	140.20	20.25	28.30	36.00
	Male	35.00	40.00	44.00	63.83	76.00	119.40	12.76	15.20	23.88	147.00	192.00	205.00	80.80	124.80	136.70	25.55	34.00	37.00
	<i>P^a</i>	0.028*			0.852			0.852			0.571			0.486			0.331		
50 – 59	Female	42.00	48.00	52.75	71.44	90.00	138.00	14.28	18.00	27.60	186.00	226.00	264.00	122.15	151.21	173.15	21.05	27.60	38.15
	Male	35.00	46.00	54.00	73.50	105.00	148.67	14.67	21.00	29.73	168.00	211.50	236.25	104.20	145.74	173.40	19.75	28.90	39.80
	<i>P^a</i>	0.288			0.613			0.613			0.152			0.344			0.714		
≥ 60	Female	46.00	52.00	55.00	72.35	102.00	164.00	14.47	20.40	32.80	179.15	207.00	265.50	114.23	130.20	155.19	24.56	31.00	39.25
	Male	33.50	36.50	43.75	69.00	88.00	159.00	13.80	17.60	31.80	177.25	193.00	231.50	119.70	132.20	159.45	22.52	32.30	40.77
	<i>P^a</i>	0.001*			0.682			0.682			0.363			0.487			0.881		

F = Female; M = Male

^a U de Mann – Whitney

* Statistically significant difference ($p < 0.05$)

Triglycerides

The reference sample for the calculation of TAG was 414 individuals (Table 4), showing a mean age of 32.90 ± 13.90 (CI 95% 31.56 to 34.24) years. In the study of the general reference sample a median of 74.69 (53.75 to

109.93) mg/dL was gained, with a median of 79.22 (50.35 to 102.53) mg/dL in females and 94.67 (57.00 to 126.63) mg/dL in males, observing statistically significant differences ($p=0.004$) between both individuals. The percentiles of this variable are shown in Table 5, and the results

of the comparisons are seen in Figure 2, panel A. When stratifying both individuals by age group to compare concentrations of triglycerides among them, significant differences in the group of 30 to 39 years were appreciated (Table 6 and Figure 4). By studying the behavior of TAG in the different age groups by gender, significant differences were observed in female individuals between groups 18-19 years and 20-29 years with the group 60 years or older ($p=0.006$ and $p=0.002$ respectively). In males, these differences were observed between the group 18-19 years with the 30-39 years group ($p=0.003$).

Figure 2: Comparison of median values of Triglycerides, Total Cholesterol and LDL-C in a sample reference by gender. Maracaibo Municipality, Zulia State, 2011.





Figure 4: Comparison of TAG and VLDL median concentrations in a reference sample by age group. Maracaibo Municipality, Zulia State, 2011.



VLDL-C


For the VLDL-C study of a total of 414 individuals of both genders were the reference sample (Table 4) with a mean age of 32.90 ± 13.90 years (CI 95% 31.56 to 34.24 years). In the general reference sample a median of 14.92 (10.75 to 21.98) mg/dL was obtained. When divided by gender, female individuals obtained a median of 14.19 (10.07 to 20.48) mg/dL while in the male group a median of 15.70 (11.40 to 25.33) mg/dL was obtained, with a statistically

significant difference $p=0.004$ (Figure 1, panel B). Percentiles for the VLDL-C concentration are given in Table 5. By stratifying according to gender and age group statistical results obtained from the VLDL concentrations analysis were the same as for the variable of triglycerides (Table 6 and Figure 4) due to the direct relationship between both lipid variables.

LDL-C

A reference sample of 422 individuals with a mean age of 33.00 ± 14.00 years (CI 95% 31.66 to 34.34 years) was used to calculate LDL concentrations (Table 4). The overall reference sample median was 110.20 (86.25 to 132.80) mg/dL. The value of the median in the female group was 109.60 (87.20 to 130.20) mg/dL; for the male group a median of 110.80 (85.60 to 134.60) mg/dL was obtained. No significant statistical difference was observed between both groups ($p=0.673$) (Figure 2, panel C). Similarly, the percentiles to determine the reference intervals for this variable by gender are expressed in Table 5. LDL-C concentration median grouped by gender and age group was obtained and shown in Table 6. No significant statistical differences were observed. The distribution of this lipid variable according to age group and gender is shown in Figure 5. When grouped by gender and age group, medians of LDL-C concentrations expressed in Table 6 were obtained, were no statistical significant differences were observed according to gender for each age group. The distribution of this lipid variable according to age group and gender is shown in Figure 5. Table 7 exhibits p values, according to different age groups for each gender.

Figure 5: Comparison of LDL-C medians of concentration in a reference sample by age groups. Maracaibo Municipality, Zulia State, 2011.



Total cholesterol

The reference sample for calculation of Total Cholesterol was 421 individuals (Table 4) with a mean age of 33.06 ± 14.08 years (CI 95% 31.71 to 34.40 years). In the overall sample a median of 176.00 (149.50 to 201.50) mg/dL was obtained. Grouped according to gender, female individuals had a median of 176.00 (151.00 to 202.00) mg/dL while men showed a median of 175.00 (147.00-200.25) mg/dL.

We found no significant differences in the analysis of both entities ($p=0.721$) (Figure 2, panel B). Reference intervals for total cholesterol according to gender are presented in Table 5. After classifying individuals according to gender and grouped by age, significant differences were observed only in the youngest group, 18-19 years, with $p=0.036$ (Figure 6). Table 8 shows p values, according to the different age groups for each gender. While analyzing each gender separately, statistically significant differences were observed between age groups located in extreme ages. In the case of females, differences were found between the age group 18 to 19 years with all groups over 40 years. As for the behavior between groups of 20-29 years and 30-39 years a difference was shown with other groups starting from the age of 50 (Figure 6). When evaluating male individuals significant differences were observed between the younger group (18-19 years) with all individuals in each subsequent age group. In addition, individuals belonging to the 20-29 years group had more difference with other groups starting from 50 years of age. The p values when comparing different age groups belonging to each gender are shown in Table 8, panel A and B.

Lipoprotein(a)

To calculate the concentrations of Lp(a), a reference sample of 329 individuals was used (Table 4) with a mean age of 33.81 ± 59 years (CI 95% 32.23 to 35.39). The overall reference sample median was 25.20 (18.40 to 33.85) mg/Dl; with a median of 25.35 (18.16 to 34.30) mg/dL for females and 24.90 (18.25 to 33.65) mg/dL for males. No significant statistical difference was shown between both groups ($p=0.571$) (Figure 1, panel C). Similarly, the percentiles to determine the reference intervals for this variable by gender are expressed in Table 5. When grouped by gender and age median concentrations of Lp(a) were obtained and expressed in Table 6 where statistically significant differences weren't observed between male and female individuals in each age group. The distribution of this lipid variable according to age group and gender is shown in Figure 7. By measuring the behavior of Lp(a) according to age group for each gender separately, there were no significant differences between different age groups for female subjects, however when assessing behavior in male individuals differences were found between the group of 18-19 years and 40-49 years group and the group of 18-19 years with the group 60 and over ($p=0.044$ and $p=0.019$ respectively).

Figure 6: Comparison of Total Cholesterol medians of concentration in a reference sample by age group. Maracaibo Municipality, Zulia State, 2011.




Table 7: P values from comparison of LDL-C concentration in individuals by gender and age group. Maracaibo Municipality, Zulia State, 2011.

Age Group (Years)	18-19	20-29	30-39	40-49	50-59	≥60
18-19	-	1.000	0.045	8.06×10^{-5} *	3.75×10^{-9} *	0.003*
20-29	1.000	-	0.073	5.81×10^{-5} *	2.92×10^{-9} *	0.006*
30-39	0.045	0.073	-	1.000	7.84×10^{-5} *	0.705
40-49	8.06×10^{-5} *	5.81×10^{-5} *	1.000	-	0.013	1.000
50-59	3.75×10^{-9} *	2.92×10^{-9} *	7.84×10^{-5} *	0.013	-	1.000
≥60	0.003*	0.006*	0.705	1.000	1.000	-

Panel B: Male

Age Group (Years)	18-19	20-29	30-39	40-49	50-59	≥60
18-19	-	0.086	2.43×10^{-4} *	0.061	6.95×10^{-7} *	6.57×10^{-7} *
20-29	0.086	-	0.220	1.000	0.001	0.004
30-39	2.43×10^{-4} *	0.220	-	1.000	1.000	1.000
40-49	0.061	1.000	1.000	-	0.543	0.591
50-59	6.95×10^{-7} *	0.001	1.000	0.543	-	1.000
≥60	6.57×10^{-6} *	0.004	1.000	0.591	1.000	-

* Statistically significant difference ($p < 0.0083$)

Table 8: p-values from comparison of Total Cholesterol concentration in individuals by gender and age group. Maracaibo Municipality, Zulia State, 2011.

Reference Intervals

HDL-C Female: 42.00 - 56.00 mg/dL.

Male: 37.00 - 50.00 mg/dL

TAG Female: 50.35 - 102.53 mg/dL.

Male: 57.00 - 126.63 mg/dL

VLDL-C Female: 10.07 - 20.48 mg/dL.

Male: 11.40 - 25.33 mg/dL.

LDL-C 86.25 - 132.80 mg/dL.

Total Cholesterol 149.50 - 201.50 mg/dL.


Lp(a) 18.40 - 33.85 mg/dL.

Table 9: Comparison of lipid profile reference intervals with others populations and commercial lab kits

Geographic Location	References	Age (Years)	LIPID PROFILE							
			Total Cholesterol		HDL-C		LDL-C		Triglycerides	
			(mg/dL)	(mmol/L)	(mg/dL)	(mmol/L)	(mg/dL)	(mmol/L)	(mg/dL)	(mmol/L)
Mexico	<i>García-Jiménez et al.</i> ¹²	Adultos	90.8 – 208.3	2.4 – 5.4	26 – 74.9	0.68 – 1.9	31.4 – 126.7	0.82 – 3.3	40.7 – 215.8	0.46 – 2.5
	<i>NOM-037-SSA2-2002</i> ²²	Adultos	<200	< 5.2	> 35	> 0.91	< 130	< 3.3	< 150	< 1.71
	<i>ENSANUT 2006</i> ²³	Adultos	162.4	4.2	40.6	1.05	121	3.1	118.1	1.5
Peru	<i>Gómez et al.</i> ³⁰	20 – 50	171 – 242	4.4 – 6.3	30 – 75	0.78 – 1.95	81.4 – 186.4	2.1- 4.8	48.– 274	0.54 – 3.1
Camerun	<i>Taga et al.</i> ³¹	20 – 30	92 – 216.8	2.38 – 5.6	30.1 – 92	0.78 – 2.38	73.1 – 175.1	1.89 – 4.53	43.7 – 185.5	0.5 -2.12
USA	<i>Heil et al.</i> ³²	Adultos	100.5 – 232	2.6 – 6	38.7 – 61.8	1 – 1.6	100.5 – 129.5	2.6 – 3.3	< 200	<2.3
	<i>Roche USA</i> *	Adultos	< 200	< 5.2	> 35	> 0.91	< 155	< 4	< 200	< 2.3
Argentina	<i>Wiener Lab</i> *	Adultos	175 – 240	4.5 – 6.2	30 – 85	0.78 – 2.2	< 140	< 3.6	35 – 165	0.39 – 1.8
Spain	<i>Spinreact</i> *	Adultos	< 200	<5.2	30 – 85	0.78 – 2.2	< 100	< 2.6	35 – 165	0.39 – 1.8
Maracaibo		Adultos	114.95-270	2.95-6.98	30-75.43	0.78- 1.94	53.64 - 185.41	1.37 - 4.78	31.88-199	0.35-2.25

* Reference values according to commercial kit

Figure 7: Comparison of Lp (a) medians of concentration in a reference sample by age group. Maracaibo Municipality, Zulia State, 2011.



Discusión

Among the various cardiovascular risk factors, dyslipidemia is one of the most important alterations involved in atherosclerosis, which is the main event behind the pathophysiology of cardiovascular diseases^{4,20,21}. In addition, the mortality rate due to cardiovascular events in the Zulia population is among the highest in the country. For these reasons, the main objective of this study was to establish lipid profile reference intervals in adult individuals of the Maracaibo Municipality, Zulia State. The infor-

mation obtained can be established as reference intervals by gender for HDL-C, TAG and VLDL-C, while for Total Cholesterol, LDL-C and Lp (a) we proposed general reference intervals since their behavior was not influenced by gender (Table 8). These findings will allow development of better diagnosis strategies that will enhance primary and secondary prevention, which will ultimately improve the quality of primary care in our locality.

Our reference population was established taking into account the recommendations of PetirClerc and Solberg²⁶, as well as the proposals of the NCEP, however, given the important influence of hyperglycemia and insulin resistance in the lipid profile³¹⁻³⁵ we decided to consider them as exclusion criteria for the selection of individuals. Based on a preliminary study from our research center we determined from a reference population that a HOMA-IR higher than 2.00 represented insulin resistance³⁵. Comparing our intervals (p2.5-p97.5) with those of other populations (Table 9), we agree with the values of HDL-C and TAG proposed by García-Jiménez et al. (Mexico) (12). Our intervals of Total Cholesterol showed a wider range than Wiener Lab (Argentina) (36) and Gómez et al. (Perú)³⁷. Also in relation to levels of LDL-C the p97.5 was one of the highest, surpassed only by that of the Peruvian population³⁸. Importantly, our p97.5 for Total Cholesterol and LDL-C significantly exceeded the cutoff points established by Roche USA, Wiener Lab and Spinreact except in HDL-C where our intervals were within the limits of the latter two laboratories.

The authors cited above agree on establishing reference intervals using the 2.5 and 97.5 percentiles as lower and upper limits respectively, however, given the alarm-

ing prevalence of metabolic endocrine disorders worldwide^{40,41} and in Venezuela, we recommend a close monitoring of patients in the percentiles 25 and 75, for which we observed interesting findings regarding the ATP III¹⁰. The 75th percentile of both Triglycerides and VLDL-C were below the cutoff points established by the ATP III to be considered normal (150 mg/dL for TAG and 30 mg/dL for VLDL-C)¹⁰. The intervals defined by such entity as borderline high level (150 - 199 mg/dL) are similar to the 90 and 97.5 percentiles of TAG observed in our reference population¹⁰. According to the epidemiological behavior of these variables (TAG and VLDL-C) by age group we observed that its concentrations remained higher in males compared to females⁴², with an accentuated statistical significant difference in individuals aged 30 to 39.

As for Total Cholesterol the p75 for reference population was 201.50 mg/dL, which is very close to the appeal of ATP III's <200 mg/dL¹⁰. Also a similar pattern of Total Cholesterol and LDL-C was found in the age groups, where both male and female individuals showed that their serum concentrations increased with age, resulting in significant differences between the extreme ages groups studied^{43,44}. In the analysis of the behavior of LDL-C levels in relation to the classification proposed by the ATP III¹⁰ it was observed that p25 is in the Optimum category (<100 mg/dL), while the p75 matches the lower value of the Borderline High category (130-159 mg/dL). Likewise, the percentiles 90 and 97.5 belong to the high category (160-189 mg/dL) without exceeding the upper limit.

Moreover, it is widely known that HDL-C is a lipid parameter that differs between genders, with higher levels for females (45). Concordant with this, our values in female individuals remained higher and the pattern held in different age groups⁴⁶⁻⁴⁸. In the independent study of each gender there were no significant differences in HDL-C concentrations going forward in age groups. By comparing the p25 of HDL-C for women and men (42.00 mg/dL and 37.00 mg/dL respectively) with the cutoff point proposed by ATP III to consider low HDL-C in an individual (50 mg/dL for women and <40 mg/dL for men) (10), we found large differences between the values of our population and the cutoff points raised by this organization, behavior that is maintained in all age groups, suggesting that age does not influence this parameter, justifying the high prevalence of hypoalphalipoproteinemia not only in our country but in our continent⁴⁹⁻⁵¹.

Although most of the literature describes that Lp(a) concentration is genetically determined and it is not significantly affected by age, gender, diet⁵² or exercise^{53,54}, our study shows that this lipoprotein concentration increases progressively with age. These findings are consistent with those obtained by Naoki Nago et al⁵⁵ who studied 1,235 men and 1,762 women over 30 years old as part of the Jichi Medical School Cohort Study conducted in Japan. In

this study we observed an increase in the concentration of Lp(a) as the subjects grew older, concluding that Lp(a) concentration is closely related to age and influenced by other factors such as alcohol consumption and gender. Similarly, in a study conducted by Jenner et al.⁵⁶ carried in 1,284 men and 1,394 women participants of the Framingham Offspring Study, a continuous and sustained rise in Lp(a) level from the age group 20 to 29 years to the group of 50 to 59 years was observed, a trend detected in both men and women that changed and decreased in the age group 60 to 69 years and in 70 or more years, very much alike to results in our study.

Despite the reported values, the rigorous process of selection of healthy patients hinders proper comparison with other populations. In the present study some characteristics of the region were not considered as eating habits or physical activity, which could influence the alteration of plasma lipid concentrations.

Conclusions

Lipid levels in our area are different from those reported by other organizations. Due to the large variability of the lipid profile, as it is characteristic of biological entities, we recommend further research both regionally and nationally, for the determination of lipid profile cutoff points and comparison thereof with the upper limit of the reference intervals, obtained in this work in the case of our population. It is necessary to establish a relationship between cardiovascular risks of each population, and, observe the reality of our situation in respect to the reported values. Likewise, the diversity of commercial test kits (even using the same analysis technique) in addition to the various sample selection methods, generates large numbers of reference intervals that prevent proper comparison of results. Therefore the standardization and consensus in the evaluation is an important premise to follow.

It should be noted that several manufacturers of laboratory reagents use as reference the arithmetic mean to determine "normal" lipid values. Due to the "abnormal" nature of biological variables and as reported in the literature we suggest that it is more convenient to use reference intervals expressed by percentiles. Likewise, we urge that clinical laboratories determine their own reference values, taking into account that eating habits, genetics and lifestyle are characteristics inherent to each population.

Acknowledgments

This work was supported by research grant No CC-0437-10-21-09-10 from Consejo de Desarrollo Científico, Tecnológico y Humanístico (CONDES) de la Universidad del Zulia, and Research grant No FZ-0058-2007 from Fundacite Zulia.

References

- Peralta M, Attie M. Enfermedad cardiovascular. Primera causa de muerte en adultos de México y el mundo. *ArchCardiolMexico*. 2007;77(2):91-93.
- Ministerio del Poder Popular para la Salud. Anuario de Mortalidad 2006. Venezuela. Available at: http://www.google.co.ve/url?sa=t&rc=j&q=2.%20ministerio%20del%20poder%20popular%20para%20la%20salud.%20anuario%20de%20mortalidad%202006.%20venezuela&source=web&cd=3&ved=0CFIQFjAC&url=http%3A%2F%2Fwww.mpps.gob.ve%2Findex.php%3Foption%3Dcom_phocadownload%26view%3Dcategory%26download%3D53%3Aanuario-2006%26id%3D11%3Aanuario-demortalidad&ei=cMm-TaxHKe_6AHWxLZNCg&usq=AFQjCNGn9Lj9vJNW_OEbnfigFpsLeLxkg
- Solano L, Velásquez E, Naddaf G, Páez M. Centro de Investigaciones en Nutrición, Facultad de Ciencias de la Salud. Universidad de Carabobo. Patron de Lípidos en Preescolares de Bajos Recursos Socio-economicos (Valencia, Venezuela). *Acta Científica Venezolana*. 2003;54:254-262.
- Caja Costaricense para el Seguro Social, Gerencia de División Médica, Dirección Técnica Servicio de Salud, Departamento de Medicina Preventiva. Guía para la Detección, Diagnóstico y Tratamiento de las Dislipidemias para el Primer Nivel de Atención. San JoseCostarrica 2004. Available at www.binasss.sa.cr/hipertension.pdf
- Dallongeville J, Fruchart JC. Postprandial Dyslipidemia: A Risk Factor for Coronary Heart Disease. *Ann NutrMetab*. 1998;42:1-11.
- Carr M, Brunzell J. Abdominal Obesity and Dyslipidemia in the Metabolic Syndrome: Importance of Type 2 Diabetes and Familial Combined Hyperlipidemia in Coronary Artery Disease Risk. *J ClinEndocrinolMetab*. 2004;89(6):2601-2607.
- Qi Q, Liang L, Doria A, Hu F, Qi, L. Genetic Predisposition to Dyslipidemia and Type 2 Diabetes Risk in Two Prospective Cohorts. *Diabetes*. 2012;61(3):745-752.
- Grundy SM. Metabolic Syndrome Pandemic. *ArteriosclerThrombVasc Biol*. 2008;28:629-636.
- Santos J. Departamento de Nutrición, Diabetes y Metabolismo Facultad de Medicina Pontificia Universidad Católica de Chile. Variación genética de lipoproteínas y lípidos plasmáticos. *RevChilCardiol*. 2010;29:88-93.
- Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. *Circulation*. 2002;106:3143.
- People's Republic of China-United States Cardiovascular and Cardiopulmonary Epidemiology Research Group. An Epidemiological Study of Cardiovascular and Cardiopulmonary Disease Risk Factors in Four Populations in the People's Republic of China. *Circulation*. 1992;85:1083-1096.
- Mesa J, Ruiz J, Ortega F, Wärnberg J, González-Lamuño D, Moreno L, Gutiérrez A, Castillo M. Aerobic physical fitness in relation to blood lipids and fasting glycaemia in adolescents: Influence of weight status. *NutrMetabCardiovascDis*. 2006;16:285-293.
- Mahley et al. Turkish Heart Study: Lipids, Lipoproteins and apolipoproteins. *J Lipid Res*. 1995;36:839-859.
- Norma Oficial Mexicana NOM-037-SSA2-2002 Para la prevención, tratamiento y control de las dislipidemias. Available at: <http://www.salud.gob.mx/unidades/cdi/nom/037ssa202.html>
- Instituto Nacional de Salud Pública. Encuesta Nacional de Salud y Nutrición. Cuernavaca, Morelos, México, 2006. Available at: www.insp.mx/ensanut/ensanut2006.pdf
- Cholestech. Standardization of Lipids Test. Technical Bulletin. 2003. Número 102.
- Das M, Saikia M, Estimation of Reference Interval of Lipid profile in Assamese Population, *Indian J ClinBiochem*. 2009;24(2):190-193.
- García-Jiménez S, Martínez M, Monroy A, Jantorena A, Sánchez M. Intervalos de referencia del perfil lipídico en trabajadores y estudiantes de la Universidad Autónoma del Estado de Morelos, México. *RevBiomed*. 2011;22:33-10.
- Pedrozo W, Bonneau G, Rascón C, Juárez M, Cardozo J. Valores de referencia y prevalencia de las alteraciones del perfil lipídico en adolescentes. *Arch Argent Pediatr*. 2010;108(2):107-115.
- Bermudez V, Paris R, Cano C, Arraiz N, Amell A, Cabrera M, Reyna N, Mengual M, Vega L, Finol F, BSc, Luti Y, Sanchez D, Sanchez W, Gonzalez J, Montes J, Rojas E, Cano J, Cano R, Velasco M, Miranda J. The Maracaibo City Metabolic Syndrome Prevalence Study: Design and Scope. *AmJTher*. 2010; 17: 288-294
- Friedewald W T, Levi R I, Fredrickson S D. Estimation of the concentration of lowdensity lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *ClinChem*1972;18(6):499-502.
- Human Gesellschaft fur Biochemica und DiagnosticambH. Lipoprotein(a), Lp(a) Prueba turbidimétrica de látex para la determinación cuantitativa de lipoproteína (a).
- Levy JC, et al. Correct homeostasis model assessment (HOMA) evaluation uses the computer program (Letter). *Diabetes Care*. 1998;21:2191-2192.
- Horn P, Feng L, Li Y. Effect of Outliers and Nonhealthy Individuals on Reference Interval Estimation. *ClinChem* 47:12 2137-2145 (2001).
- Solberg, H. Approved Recommendation (1986) on the theory of reference values. Part 1. The concept of reference values. *ClinicaChimica Acta*. 1987;165:111-118.
- PetitClerc, C and Solberg H. Approved recommendation (1987) on the theory of reference values. Part 2. Selection of individuals for the production of reference values. *J ClinChemClinBiochem*. 1987;25:639-644.
- Queraltó J, Amoja F, Cortes M, Domenech M, Fuemes J, L1agoslera M, Ordonez J, Ramon F, Rojo I. Sociedad Española de Química Clínica, Comité Científico, Comisión Valores de Referencia. Concepto de valores de referencia en Química Clínica. *Química Clínica*. 1983;2(1):39-41.
- Queraltó J, Amoja F, Cortes M, Domenech M, Fuemes J, L1agoslera M, Ordonez J, Ramon F, Rojo I. Sociedad Española de Química Clínica, Comité Científico, Comisión Valores de Referencia. Variaciones analíticas y extra-analíticas en la producción de los Valores de Referencia. *Química Clínica*. 1984;3(1):43-50.
- Sara García-Jiménez. Intervalos de referencia del perfil lipídico en trabajadores y estudiantes de la Universidad Autónoma del Estado de Morelos, México. *RevBiomed* 2011; 22:3-10.
- Tukey JW. *Exploratory data analysis*. Reading, MA: Addison-Wesley. 1977:506pp.
- American Heart Association Heart and Stroke Statistics Committee. 2008. Available at: http://www.americanheart.org/downloadable/heart/1200078608862HS_Stats%202008.final.pdf
- Grundy SM, Balady GJ, Criqui MH, Fletcher G, Greenland P, Hiratzka LF, et al. Primary prevention of coronary heart disease: guidance from Framingham: a statement for healthcare professionals from the AHA Task Force on Risk Reduction. American Heart Association. *Circulation* 1998;97:1876-1887.
- Bettina Mittendorfer, Bruce W. Patterson, Samuel Klein and Labros S. Sidossis. VLDL-triglyceride kinetics during hyperglycemia-hyperinsulinemia: effects of sex and obesity. *Am J PhysiolEndocrinolMetab*2003;284:E708-E715.
- Aarsland A, Chinkes D, and Wolfe RR. Contributions of the novo synthesis of fatty acids to total VLDL-triglyceride secretion during prolong hyperglycemia/hyperinsulinemia in normal man. *J ClinInvest*. 1996; 98: 2008-2017.
- Bermúdez V. Estimación de los valores de referencia de sensibilidad insulínica por Homeostasis ModelAssessment (HOMA) en individuos adultos del municipio Maracaibo, estado Zulia. *Diabetes Internacional* 2010;11(1):19.
- Wiener Laboratorios S.A.I.C 2000. Rosario- Argentina. <http://www.wiener-lab.com.ar>
- Gomez TPJC, Bustinza LE, Huarachi A. Valores de referencia de algunas pruebas bioquímicas y hematológicas en personas adultas sanas del Hospital Central de la Fuerza Aerea del Perú 2000-2001. *RevMex Patol Clin* 2003;50:41-9.
- Taga I, Kouemeli N, L gogang- Yokeu J. Valeurs de références des lipidessériquechez de jeuneshommescamerounais. *Ann BiolClin* 2004;41:33-8.
- Heil W, Koberstein R, Zawta B. Reference ranges for adults and children. Roche Diagnostic USA. 2004.
- Goff D. Jr; Bertoni A; Kramer H; Bonds D; Blumenthal R; Tsai M; Psaty B. Dyslipidemia Prevalence, Treatment, and Control in the Multi-Ethnic Study of Atherosclerosis (MESA): Gender, Ethnicity, and Coronary Artery Calcium. *Circulation*. 2006;113:647-656.
- Ogden C, Yanovski S, Carroll M, Flegal K. The Epidemiology of Obesity. *Gastroenterology*. 2007;132:2087-2102.
- Carroll MD, Lacher DA, Sorlie PD, Cleeman JI, Gordon DJ, WOlz M, Grundy SM, Johnson CL. Trends in serum lipids and lipoproteins of adults, 1960-2002. *JAMA*. 2005;294:1773-1781.
- The Lipid Research Clinics Program Epidemiology Committee. Plasma lipid distributions in selected North American populations: The Lipid Reaserch Clinics Program Prevalence Study. *Circulation*. 1979;60:427-439.

44. Verschuren WM, Boerma GJ, Kromhout D. Total and HDL cholesterol in The Netherlands: 1987–1992. Levels and changes over time in relation to age, gender and educational level. *Int J Epidemiol*. 1994;23:948–956.
45. Schaefer EA, Lamon-Fava S, Cohn SD, Schaefer MM, Ordovas JM, Castelli WP, F. WPW. Effects of age, gender, and menopausal status on plasma low density lipoprotein cholesterol and apolipoprotein B levels in the Framingham Offspring Study. *J Lipid Res*. 1994;35:779–792.
46. Heiss G, Tamir I, Davis CE, Tyroler HA, Rifkind BM, Schonfeld G, Jacobs D, Frantz ID. Lipoprotein-cholesterol distributions in selected North America populations: the Lipid Research Clinics Program Prevalence Study. *Circulation*. 1980;61:302-315.
47. Heitmann BL. The effects of gender and age on associations between blood lipid levels and obesity in Danish men and women aged 35-65 years. *J ClinEpidemiol*. 1992;45:693-702.
48. Abbott RD, Garrison RJ, Wilson PW, Epstein FH, Castelli WP, Feinleib M, LaRue C. Joint distribution of lipoprotein cholesterol classes: the Framingham Study. *Arteriosclerosis*. 1983;3:260-272.
49. Aguilar-Salinas C et al. High Prevalence of Low HDL cholesterol concentrations and mixed hyperlipidemia in a Mexican nationwide survey. *JLipid Res*. 2001;42:1298-1307.
50. Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults. *JAMA* 2002; 287:356–359.
51. Bautista L, et al. Prevalence and impact of cardiovascular risk factors in Bucaramanga, Colombia: results from the Country wide Integrated Noncommunicable Disease Intervention Programme (CINDI/CARMEN) baseline survey. *Eur J CardiovascPrevRehabil* 2006;13:769–775.
52. Krause M, Sonntag B, Klamroth R. Lipoprotein(a) and other prothrombotic risk factors in Caucasian women with unexplained recurrent miscarriage. Results of a multicentre case-control study. *ThrombHaemost*2005;93(5):867-71.
53. Hubinger L, Mackinnon LT. The effect of endurance training on lipoprotein(a) [Lp(a)] levels in middle-aged males. *Med Sci Sports Exerc*1996;28(6):757-764.
54. Hubinger L, Mackinnon LT, Lepre F. Lipoprotein(a) [Lp(a)] levels in middle-aged male runners and sedentary controls. *Med Sci Sports Exerc*1995;27(4):490-496.
55. Nago N, Kayaba K, Hiraoka J, Matsuo H, Goto T, Kario K, Tsutsumi A, Nakamura Y, Igarashi M. Lipoprotein(a) Levels in the Japanese Population: Influence of Age and Sex, and Relation to Atherosclerotic Risk Factors: The Jichi Medical School Cohort Study. *Am J Epidemiol*1995;141:815–21.
56. JL Jenner, JM Ordovas, S Lamon-Fava, MM Schaefer, PW Wilson, WP Castelli and EJ Schaefer. Effects of age, sex, and menopausal status on plasma lipoprotein(a) levels. The Framingham Offspring Study. *Circulation* 1993;87;1135-1141.



Experto en Mercadeo Farmacéutico
Coordinador Académico: Prof. Andrés Reyes Polanco

ESTUDIA en ESPAÑA
desde VENEZUELA
DIRIGIDOS A PROFESIONALES UNIVERSITARIOS

Título propio de la Universidad de Alcalá, con una carga académica de 30 créditos ECTS (750 horas lectivas), aprobado por la Comisión de Postgrado y supervisado por el Instituto de Calidad Español - ICE.

El alumno es jurídica y académicamente matriculado y estudiando en la Universidad de Alcalá.

Culminados con éxitos los estudios, el alumno recibe el título de Experto en... por la Universidad de Alcalá, firmado por el Rector y legalizado con la Apostilla de La Haya, tiene valor académico universal.

En ningún caso está cursando una "especialidad venezolana", ni obtiene título alguno de la República Bolivariana de Venezuela.

Información: (0212) 881.19.07 / css212518@gmail.com